Neuroscience Letters, 98 (1989) 356–360
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abó, V., Viera, L., Silveira, R. and Dajas, F., Effects of local inhibition of locus coeruleus acetylcholinesterase by fasciculin in rats, 253
Akaoka, H., Vitte, P.A., Lestage, P., Saunier, C.F., Bobillier, P. and Chouvet, G., Apomorphine-induced inhibition of substantia nigra dopamine neurons: effects of unilateral injection through the internal carotid artery, 205
Åkerman, K.E.O, see Holopainen, I., 57
Alho, K., see Näätänen, R., 217
Amato, G., see Salerno, M.T., 299
Anderson, P.N., see Woodham, P., 8
Arai, O., Taniguchi, I. and Saito, N., Correlation between the size of song control nuclei and plumage color change in orange bishop birds, 144
Argiolas, A., see Melis, M.R., 351
Arnold, A.P., see Lee, J.H., 79
Avoli, M., see Hwa, G.G.C., 189

Baimbridge, K.G., see Obenaus, A., 172
Barbanti, G., see Maggi, C.A., 317
Batten, T.F.C. and Cambre, M.L., Calcitonin gene-related peptide-like immunoreactive fibres innervating the hypothalamic inferior lobes of teleost fishes, 1
Bennett, G.J., see Sugimoto, T., 139
Bernstein, H.-G., Reichenbach, A., Kirschke, H. and Wiederanders, B., Cell type-specific distribution of cathepsin B and D immunoreactivity within the rabbit retina, 135
Biesold, D., Inamai, O., Sato, A. and Sato, Y., Stimulation of the nucleus basalis of Meynert increases cerebral cortical blood flow in rats, 39
Bobillier, P., see Akaoka, H., 205
Brailoiu, E., see Brănişteanu, D.D., 96
Brănişteanu, D.D., Brănişteanu, D.D.D., Covic, A., Brailoiu, E., Serban, D.N. and Haulica, I.D., Adenosine effects upon the spontaneous quantal transmitter release at the frog neuromuscular junction in the presence of protein kinase C-blocking and -activating agents, 96
Brănişteanu, D.D.D., see Brănişteanu, D.D., 96
Bures, J. and Buresova, O., Conditioned taste aversion to injected flavor: differential effect of anesthesia on the formation of the gustatory trace and on its association with poisoning in rats, 305
Buresova, O., see Bures, J., 305

Cambre, M.L., see Batten, T.F.C., 1
Caorsi, C.E. and González, C.B., Antibodies against specific membrane proteins of neurosecretory granules isolated from bovine neural lobes, 241
Castro, A.J., see Sørensen, J.Chr., 33
Chouvet, G., see Akaoka, H., 205
Compton, R.P., see Hood, W.F., 91
Cortimiglia, R., see Salerno, M.T., 299
Coulter, D.A., Huguenard, J.R. and Prince, D.A., Specific petit mal anticonvulsants reduce calcium current in thalamic neurons, 74
Covic, A., see Brănişteanu, D.D., 96
Crescimanno, G., see Salerno, M.T., 299
Cunningham, J.R., see Neal, M.J., 29

Dajas, F., see Abó, V., 253
Danguir, J. and De Saint-Hilaire Kafi, S., Reversal of desipramine-induced suppression of paradoxical sleep by a long-acting somatostatin analogue (octreotide) in rats, 154
De Carlos, J.A., see López-Mascaraque, L., 19
De Saint-Hilaire Kafi, S., see Danguir, J., 154
Del Bianco, E., see Maggi, C.A., 317
Demeneix, B.A., see Trouslard, J., 234
Di Paolo, T., see Lévesque, D., 345
Dingledine, R., see Traynelis, S.F., 194
Drejer, J., Sheardown, M., Nielsen, E.Ø. and Honoré, T., Glycine reverses the effect of HA-966 on NMDA responses in cultured rat cortical neurons and in chick retina, 333
Drew, K.L., see O'Connor, W.T., 211

Ebinger, G., see Herregodts, P., 321
Enkvist, M.O.K., see Holopainen, I., 57

Fabian, R.H., see Hulsebosch, C.E., 13
Feltz, P., see Trouslard, J., 234
Ferraro, G., see Sabatino, M., 285
Ferrer, I., see Garcia-Verdugo, J.M., 247
Flint Beal, M., see Kowal, N.W., 118
Fujiwara, T. and Nagakuro, C., Three-dimensional structure of the presynaptic nerve ending in the ciliary ganglion of the chick embryo: a scanning electron microscopic study, 125

Gagnon, S., see Lévesque, D., 345

0304-3940/89/$ 03.50 © 1989 Elsevier Scientific Publishers Ireland Ltd.

Neuroscience Letters, 98 (1989) 356–360
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abó, V., Viera, L., Silveira, R. and Dajas, F., Effects of local inhibition of locus coeruleus acetylcholinesterase by fasciculin in rats, 253
Akaoka, H., Vitte, P.A., Lestage, P., Saunier, C.F., Bobillier, P. and Chouvet, G., Apomorphine-induced inhibition of substantia nigra dopamine neurons: effects of unilateral injection through the internal carotid artery, 205
Åkerman, K.E.O, see Holopainen, I., 57
Alho, K., see Näätänen, R., 217
Amato, G., see Salerno, M.T., 299
Anderson, P.N., see Woodham, P., 8
Arai, O., Taniguchi, I. and Saito, N., Correlation between the size of song control nuclei and plumage color change in orange bishop birds, 144
Argiolas, A., see Melis, M.R., 351
Arnold, A.P., see Lee, J.H., 79
Avoli, M., see Hwa, G.G.C., 189

Baimbridge, K.G., see Obenaus, A., 172
Barbanti, G., see Maggi, C.A., 317
Batten, T.F.C. and Cambre, M.L., Calcitonin gene-related peptide-like immunoreactive fibres innervating the hypothalamic inferior lobes of teleost fishes, 1
Bennett, G.J., see Sugimoto, T., 139
Bernstein, H.-G., Reichenbach, A., Kirschke, H. and Wiederanders, B., Cell type-specific distribution of cathepsin B and D immunoreactivity within the rabbit retina, 135
Biesold, D., Inamai, O., Sato, A. and Sato, Y., Stimulation of the nucleus basalis of Meynert increases cerebral cortical blood flow in rats, 39
Bobillier, P., see Akaoka, H., 205
Brailoiu, E., see Brănişteanu, D.D., 96
Brănişteanu, D.D., Brănişteanu, D.D.D., Covic, A., Brailoiu, E., Serban, D.N. and Haulica, I.D., Adenosine effects upon the spontaneous quantal transmitter release at the frog neuromuscular junction in the presence of protein kinase C-blocking and -activating agents, 96
Brănişteanu, D.D.D., see Brănişteanu, D.D., 96
Bures, J. and Buresova, O., Conditioned taste aversion to injected flavor: differential effect of anesthesia on the formation of the gustatory trace and on its association with poisoning in rats, 305
Buresova, O., see Bures, J., 305

Cambre, M.L., see Batten, T.F.C., 1
Caorsi, C.E. and González, C.B., Antibodies against specific membrane proteins of neurosecretory granules isolated from bovine neural lobes, 241
Castro, A.J., see Sørensen, J.Chr., 33
Chouvet, G., see Akaoka, H., 205
Compton, R.P., see Hood, W.F., 91
Cortimiglia, R., see Salerno, M.T., 299
Coulter, D.A., Huguenard, J.R. and Prince, D.A., Specific petit mal anticonvulsants reduce calcium current in thalamic neurons, 74
Covic, A., see Brănişteanu, D.D., 96
Crescimanno, G., see Salerno, M.T., 299
Cunningham, J.R., see Neal, M.J., 29

Dajas, F., see Abó, V., 253
Danguir, J. and De Saint-Hilaire Kafi, S., Reversal of desipramine-induced suppression of paradoxical sleep by a long-acting somatostatin analogue (octreotide) in rats, 154
De Carlos, J.A., see López-Mascaraque, L., 19
De Saint-Hilaire Kafi, S., see Danguir, J., 154
Del Bianco, E., see Maggi, C.A., 317
Demeneix, B.A., see Trouslard, J., 234
Di Paolo, T., see Lévesque, D., 345
Dingledine, R., see Traynelis, S.F., 194
Drejer, J., Sheardown, M., Nielsen, E.Ø. and Honoré, T., Glycine reverses the effect of HA-966 on NMDA responses in cultured rat cortical neurons and in chick retina, 333
Drew, K.L., see O'Connor, W.T., 211

Ebinger, G., see Herregodts, P., 321
Enkvist, M.O.K., see Holopainen, I., 57

Fabian, R.H., see Hulsebosch, C.E., 13
Feltz, P., see Trouslard, J., 234
Ferraro, G., see Sabatino, M., 285
Ferrer, I., see Garcia-Verdugo, J.M., 247
Flint Beal, M., see Kowal, N.W., 118
Fujiwara, T. and Nagakuro, C., Three-dimensional structure of the presynaptic nerve ending in the ciliary ganglion of the chick embryo: a scanning electron microscopic study, 125

Gagnon, S., see Lévesque, D., 345

0304-3940/89/$ 03.50 © 1989 Elsevier Scientific Publishers Ireland Ltd.

Garcia-Verdugo, J.M., Llahi, S., Ferrer, I. and Lopez-Garcia, C., Postnatal neurogenesis in the olfactory bulbs of a lizard. A tritiated thymidine autoradiographic study, 247
Geppetti, P., see Maggi, C.A., 317
Gessa, G.L., see Melis, M.R., 351
González, C.B., see, Caorsi, C.E., 241
Goodlett, C.R., see Hamre, K.M., 264
Grafe, P., see Spuler, A., 280
Gravante, G., see Sabatino, M., 285
Grewal, A., see Mathers, D.A., 229
Guiramand, J., Nourigat, A., Sassetti, I. and Recasens, M., K^+ differentially affects the excitatory amino acids- and carbachol-elicited inositol phosphate formation in rat brain synaptoneurosomes, 222

Hamre, K.M., Hyman, B.T., Goodlett, C.R., West, J.R. and Van Hoesen, G.W., Alz-50 immunoreactivity in the neonatal rat: changes in development and co-distribution with MAP-2 immunoreactivity, 264
Hartschuh, W. and Weihe, E., Pancreastatin-like immunoreactivity in epidermal Merkel cells of pig and man, 258
Haulica, I.D., see Brănişteanu, D.D., 96
Hayashi, S., see Tanaka, Y., 179
Hepp-Reymond, M.-C., see Wannier, T.M.J., 63
Herregodts, P., Michotte, Y. and Ebinger, G., Regional differences in the distribution of norephinephrine and epinephrine in human cerebral cortex: a neurochemical study using HPLC and electrochemical detection, 321
Holopainen, I., Enkvist, M.O.K. and Åkerman, K.E.O., Glutamate receptor agonists increase intracellular Ca^{2+} independently of voltage-gated Ca^{2+} channels in rat cerebellar granule cells, 57
Honegger, C.G., Krenger, W. and Langemann, H., Measurement of free radical scavengers in the spinal cord of rats with experimental autoimmune encephalomyelitis, 327
Honoré, T., see Drejer, J., 333
Hood, W.F., Compton, R.P. and Monahan, J.B., D-Cycloserine: a ligand for the *N*-methyl-D-aspartate coupled glycine receptor has partial agonist characteristics, 91
Huesgen, A., see Vollrath, L., 297
Huguenard, J.R., see Coulter, D.A., 74
Hulsebosch, C.E. and Fabian, R.H., Penetration of IgGs into the neuraxis of the neonatal rat, 13
Hwa, G.G.C. and Avioli, M., NMDA receptor antagonists CPP and MK-801 partially suppress the epileptiform discharges induced by the convulsant drug bicuculline in the rat neocortex, 189
Hyman, B.T., see Hamre, K.M., 264

Igarashi, K., see Onodera, H., 101
Inamai, O., see Biesold, D., 39
Ishibashi, Y., see Mizuno, A., 272
Iwai, C., see Shioda, S., 25

Johnson, R.F., Smale, L., Moore, R.Y. and Morin, L.P., Paraventricular nucleus efferents mediating photoperiodism in male golden hamsters, 85
Jordan, C.L., see Lee, J.H., 79

Kajander, K.C., see Sugimoto, T., 139
Kato, M., Chronically isolated lumbar half spinal cord, produced by hemisection and longitudinal myelotomy, generates locomotor activities of the ipsilateral hindlimb of the cat, 149
Kawamura, M., see Minami, M., 105
Keller, F., see Lorez, H., 339
Kirschke, H., see Bernstein, H.-G., 135
Kiyota, Y., see Onodera, H., 101
Kogure, K., see Onodera, H., 101
Kowal, N.W. and Flint Beal, M., Galanin-like immunoreactivity is present in human substantia innominata and in senile plaques in Alzheimer's disease, 118
Krenger, W., see Honegger, C.G., 327
Kriegstein, A.R., see Shen, J.M., 184
Kuraishi, Y., see Minami, M., 105
Kurosawa, M., Sato, A. and Sato, Y., Stimulation of the nucleus basalis of Meynert increases acetylcholine release in the cerebral cortex in rats, 45

La Grutta, G., see Sabatino, M., 285
La Grutta, V., see Sabatino, M., 285
Langemann, H., see Honegger, C.G., 327
Lee, J.H., Jordan, C.L. and Arnold, A.P., Critical period for androgenic regulation of soma size of sexually dimorphic motoneurons in rat lumbar spinal cord, 79
Leiderer, P., see Vollrath, L., 297
Lestage, P, see Akaoka, H., 205
Li, Y.Q., Rao, Z.R. and Shi, J.W., Serotoninergic projections from the midbrain periaqueductal gray to the nucleus accumbens in the rat, 276
Llahi, S., see Garcia-Verdugo, J.M., 247

Loeffler, J.Ph., see Trouslard, J., 234
Lopez-Garcia, C., see Garcia-Verdugo, J.M., 247
Lorez, H., Keller, F., Ruess, G. and Otten, U., Nerve growth factor increases in adult rat brain after hypoxic injury, 339
Lévesque, D., Gagnon, S. and Di Paolo, T., Striatal D1 dopamine receptor density fluctuates during the rat estrous cycle, 345
López-Mascaraque, L., Villalba, R.M. and De Carlos, J.A., Vasocactive intestinal polypeptide-immunoreactive neurons in the main olfactory bulb of the hedgehog (*Erinaceus europaeus*), 19

Maggi, C.A., Santicioli, P., Del Bianco, E., Geppetti, P., Barbanti, G., Turini, D. and Meli, A., Release of VIP- but not CGRP-like immunoreactivity by capsaicin from the human isolated small intestine, 317
Maier, M.A., see Wannier, T.M.J., 63
Manz, B., see Vollrath, L., 297
Masu, Y., see Minami, M., 105
Mathers, D.A., Grewal, A. and Wang, Y., Membrane channels activated by taurine in cultured mouse spinal cord neurons, 229
Matsuda, M., see Mizuno, A., 272
Matthies, H., see Reymann, K.G., 166
Meli, A., see Maggi, C.A., 317
Melis, M.R., Argiolas, A. and Gessa, G.L., Apomorphine increases plasma oxytocin concentration in male rats. 351
Michotte, Y., see Herregodts, P., 321
Minami, M., Kuraishi, Y., Kawamura, M., Yamaguchi, T., Masu, Y., Nakanishi, S. and Satoh, M., Enhancement of preprotachykinin A gene expression by adjuvant-induced inflammation in the rat spinal cord: possible involvement of substance P-containing spinal neurons in nociception, 105
Misgeld, U. and Swandulla, D., Quisqualate receptor-mediated rhythmic bursting of rat hypothalamic neurons in dissociated cell culture, 291
Mizobuchi, T., see Mizuno, A., 272
Mizuno, A., Mizobuchi, T., Ishibashi, Y. and Matsuda, M., C-Fos mRNA induction under vitamin B_6 antagonist-induced seizure, 272
Mody, I., see Obenaus, A., 172
Monahan, J.B., see Hood, W.F., 91
Moore, R.Y., see Johnson, R.F., 85
Morin, L.P., see Johnson, R.F., 85

Nadim, W., see Woodham, P., 8
Nagakuro, C., see Fujiwara, T., 125
Nagaoka, A., see Onodera, H., 101
Nakai, Y., see Shioda, S., 25
Nakanishi, S., see Minami, M., 105
Neal, M.J., Cunningham, J.R., Shah, M.A. and Yazulla, S., Immunocytochemical evidence that vigabatrin in rats causes GABA accumulation in glial cells of the retina, 29
Nielsen, E.Ø., see Drejer, J., 333
Nourigat, A., see Guiramand, J., 222
Näätänen, R., Paavilainen, P., Alho, K., Reinikainen, K. and Sams, M., Do event-related potentials reveal the mechanism of the auditory sensory memory in the human brain?, 217

O'Connor, W.T., Drew, K.L. and Ungerstedt, U., Differences in dopamine release and metabolism in rat striatal subregions following acute clozapine using in vivo microdialysis, 211
Obenaus, A., Mody, I. and Baimbridge, K.G., Dantrolene-Na (Dantrium) blocks induction of long-term potentiation in hippocampal slices, 172
Ono, Y., see Onodera, H., 101
Onodera, H., Kogure, K., Ono, Y., Igarashi, K., Kiyota, Y. and Nagaoka, A., Proto-oncogene c-fos is transiently induced in the rat cerebral cortex after forebrain ischemia, 101
Otten, U., see Lorez, H., 339

Paavilainen, P., see Näätänen, R., 217
Palovcik, R.A., see Reid, S.A., 200
Pastuszko, A., Yee, D.K. and Wilson, D.F., Modulation of catecholamine metabolism in synaptosomes by a neuroregulatory factor from mammalian brain, 111
Pevet, P., see Skene, D.J., 69
Pollow, K., see Vollrath, L., 297
Prince, D.A., see Coulter, D.A., 74

Rao, Z.R., see Li, Y.Q., 276
Rasmusson, D.D., The projection pattern of forepaw nerves to the cuneate nucleus of the raccoon, 129
Recasens, M., see Guiramand, J., 222
Reichenbach, A., see Bernstein, H.-G., 135
Reid, S.A. and Palovcik, R.A., Spontaneous epileptiform discharges in isolated human cortical slices from epileptic patients, 200
Reinikainen, K., see Näätänen, R., 217
Reymann, K.G. and Matthies, H., 2-Amino-4-phosphonobutyrate selectively eliminates late

phases of long-term potentiation in rat hippocampus, 166
Ruess, G., see Lorez, H., 339

Sabatino, M., Gravante, G., Ferraro, G., Vella, N., La Grutta, G. and La Grutta, V., Striatonigral suppression of focal hippocampal epilepsy, 285
Saito, H., see Tanaka, J., 51
Saito, N., see Arai, O., 144
Sakurai, M., see Tanaka, Y., 179
Salerno, M.T., Cortimiglia, R., Crescimanno, G. and Amato, G., Effect of claustrum activation on the spontaneous unitary activity of frontal eye field neurons in the cat, 299
Sams, M., see Näätänen, R., 217
Santicioli, P., see Maggi, C.A., 317
Sassetti, I., see Guiramand, J., 222
Sato, A., see Biesold, D., 39
Sato, A., see Kurosawa, M., 45
Sato, Y., see Bieslold, D., 39
Sato, Y., see Kurosawa, M., 45
Satoh, M., see Minami, M., 105
Saunier, C.F., see Akaoka, H., 205
Schenkel, E. and Siegel, J.M., REM sleep without atonia after lesions of the medial medulla, 159
Schulz, D.W. and Zigmond, R.E., Neuronal bungarotoxin blocks the nicotinic stimulation of endogenous dopamine release from rat striatum, 310
Seidel, A., see Vollrath, L., 297
Serban, D.N., see Brănişteanu, D.D., 96
Shah, M.A., see Neal, M.J., 29
Sheardown, M., see Drejer, J., 333
Shen, J.M. and Kriegstein, A.R., The development of bicuculline-induced epileptiform discharges in embryonic turtle cortex, 184
Shi, J.W., see Li, Y.Q., 276
Shioda, S., Nakai, Y., Iwai, C. and Sunayama, H., Co-existence of TRH with mesotocin in the same axon terminals of the bullfrog pars nervos as revealed by double labeling immunocytochemistry, 25
Siegel, J.M., see Schenkel, E., 159
Silveira, R., see Abó, V., 253
Skene, D.J., Vivien-Roels, B. and Pevet, P., Pineal 5-methoxytryptophol rhythms in the box turtle: effect of photoperiod and environmental temperature, 69
Smale, L., see Johnson, R.F., 85
Sørensen, J.Chr., Zimmer, J. and Castro, A.J., Fetal cortical transplants reduce the thalamic atrophy induced by frontal cortical lesions in newborn rats, 33
Spuler, A. and Grafe, P., Adenosine, 'pertussis-sensitive' G-proteins, and K^+ conductance in central mammalian neurones under energy deprivation, 280
Sugimoto, T., Bennett, G.J. and Kajander, K.C., Strychine-enhanced transsynaptic degeneration of dorsal horn neurons in rats with an experimental painful peripheral neuropathy, 139
Sunayama, H., see Shioda, S., 25
Swandulla, D., see Misgeld, U., 291

Tanaka, J., Saito, H. and Yagyu, K., Impaired responsiveness of paraventricular neurosecretory neurons to osmotic stimulation in rats after local anesthesia of the subfornical organ, 51
Tanaka, Y., Sakurai, M. and Hayashi, S., Effect of scopolamine and HP 029, a cholinesterase inhibitor, on long-term potentiation in hippocampal slices of the guinea pig, 179
Taniguchi, I., see Arai, O., 144
Traynelis, S.F. and Dingledine, R., Modification of potassium-induced interictal bursts and electrographic seizures by divalent cations, 194
Trouslard, J., Loeffler, J.Ph., Demeneix, B.A. and Feltz, P., Thyrotropin-releasing hormone stimulates porcine melanotrope cells in primary culture, 234
Turini, M., see Maggi, C.A., 317
Turmaine, M., see Woodham, P., 8

Ungerstedt, U., see O'Connor, W.T., 211
Van Hoesen, G.W., see Hamre, K.M., 264
Vella, N., see Sabatino, M., 285
Viera, L., see Abó, V., 253
Villalba, R.M., see López-Mascaraque, L., 19
Vitte, P.A., see Akaoka, H., 205
Vivien-Roels, B., see Skene, D.J., 69
Vollrath, L., Seidel, A., Huesgen, A., Manz, B., Pollow, K. and Leiderer, P., One millisecond of light suffices to suppress nighttime pineal melatonin synthesis in rats, 297

Wang, Y., see Mathers, D.A., 229
Wannier, T.M.J., Maier, M.A. and Hepp-Reymond, M.-C., Responses of motor cortex neurons to visual stimulation in the alert monkey, 63
Weihe, E., see Hartschuh, W., 258
West, J.R., see Hamre, K.M., 264
Wiederanders, B., see Bernstein, H.-G., 135

Wilson, D.F., see Pastuszko, A., 111
Woodham, P., Anderson, P.N., Nadim, W. and Turmaine, M., Satellite cells surrounding axotomised rat dorsal root ganglion cells increase expression of GFAP-like protein, 8
Yagyu, K., see Tanaka, J., 51
Yamaguchi, T., see Minami, M., 105
Yazulla, S., see Neal, M.J., 29
Yee, D.K., see Pastuszko, A., 111
Zigmond, R.E., see Shulz, D.W., 310
Zimmer, J., see Sørensen, J.Chr., 33

CONTENTS *(continued from back cover)*

Measurement of free radical scavengers in the spinal cord of rats with experimental autoimmune encephalomyelitis (NSL 05972)
C.G. Honegger, W. Krenger and H. Langemann (Switzerland) 327
Glycine reverses the effect of HA-966 on NMDA responses in cultured rat cortical neurons and in chick retina (NSL 05973)
J. Drejer, M. Sheardown, E.Ø. Nielsen and T. Honoré (Denmark) 333
Nerve growth factor increases in adult rat brain after hypoxic injury (NSL 05974)
H. Lorez, F. Keller, G. Ruess and U. Otten (Switzerland) 339
Striatal D_1 dopamine receptor density fluctuates during the rat estrous cycle (NSL 05976)
D. Lévesque, S. Gagnon and T. Di Paolo (Canada) 345
Apomorphine increases plasma oxytocin concentration in male rats (NSL 05958)
M.R. Melis, A. Argiolas and G.L. Gessa (Italy) 351